


Problems - Chapter 13 (without solutions)

1) The phase diagram for a pure chemical substance is given below, and may be used to answer the following questions.



a) What are the values for p and T at the triple point of the substance?

b) What are the values for p and T at the critical point of the substance?

c) Give the temperature and pressure corresponding to the normal freezing point, the normal boiling point, and the normal sublimation point for the substance. If a normal point for any of these phase transitions does not exist, explain why the point does not exist.

2) Define the following terms: a) solution, b) miscible, c) hydration, d) percent by mass (solute), e) colligative property, f) hypotonic.

3) The following question concerns mixing of liquids.

a) Consider mixing two liquids where mixing is exothermic ($\Delta H_{\text{soln}} < 0$). Would you expect a solution to form (yes/maybe/no)? Justify your answer.

b) Consider mixing two liquids where mixing is endothermic ($\Delta H_{\text{soln}} > 0$). Would you expect a solution to form (yes/maybe/no)? Justify your answer.

4) Explain the variation in the solubility in water of the listed alcohols:

<u>Compound</u>	<u>Solubility (g/100 g H₂O) at T = 20. °C</u>
CH ₃ OH	∞
CH ₃ CH ₂ OH	∞
CH ₃ CH ₂ CH ₂ OH	∞
CH ₃ CH ₂ CH ₂ CH ₂ OH	9.0
CH ₃ CH ₂ CH ₂ CH ₂ CH ₂ OH	2.7


∞ means the alcohol and water are completely miscible

5) Which of the following liquids is expected to be miscible with water – methanol (CH₃OH), acetone (CH₃COCH₃), cyclohexane (C₆H₁₂)?


6) What does the statement like dissolves like mean with respect to solution formation?

7) For each compound would you expect greater solubility in water (H₂O) or in hexane (C₆H₁₄)? Indicate the kinds of intermolecular forces that would occur between the solute and the solvent in which the molecule is more soluble.


a. toluene




b. sucrose
(table sugar)



c. isobutene



d. ethylene glycol



8) Calculate the molality of each of the following aqueous solutions.

a) 2.55 M NaCl solution, density = D = 1.08 g/mL.

b) 45.2 percent by mass KBr.

9) The density of an aqueous solution containing 15.0 percent ethyl alcohol (EtOH = $\text{CH}_3\text{CH}_2\text{OH}$) by mass is $D = 0.984 \text{ g/mL}$.

a) Calculate the molality of this solution.

b) Calculate the molarity of the solution

10) A solution is 14.8% by mass hexane (C_6H_{14} , $\text{MW} = 86.2 \text{ g/mol}$) in carbon tetrachloride (CCl_4 , $\text{MW} = 153.8 \text{ g/mol}$). What are the molality and the mole fraction of hexane in the solution?

11) An aqueous KNO_3 solution is made using 72.5 g of KNO_3 dissolved to a total solution volume of 2.00 L. Calculate the molarity, molality, and mass percent KNO_3 in the solution. Assume a density of 1.050 g/mL for the solution.

12) A stock solution is prepared by dissolving 18.21 g of sodium acetate ($\text{NaC}_2\text{H}_3\text{O}_2$, $\text{MW} = 82.03 \text{ g/mol}$) in water. The final volume of the solution is $V = 250.0 \text{ mL}$.

a) What is the molarity of sodium acetate in the solution?

b) How many mL of the stock solution of sodium acetate are needed to prepare 100.0 mL of a 0.1000 mol/L sodium acetate solution?


13) The Henry's law constant for methyl bromide (CH_3Br , $\text{MW} = 94.9 \text{ g/mol}$) in water is $0.159 \text{ mol/L}\cdot\text{atm}$ at $T = 25. \text{ }^\circ\text{C}$. How many grams of methyl bromide will dissolve in 1.000 L of water at this temperature when the pressure of methyl bromide above the water is $p = 320. \text{ torr}$?

14) The following question concerns ideal solutions.

a) How do we define an ideal solution of two gases?

b) How do we define an ideal solution of two volatile liquids?

15) The diagram below is for a solution of two volatile liquids A and B, and shows the partial pressure of A, the partial pressure of B, and the total pressure above a solution as a function of X_A , the mole fraction of A in the solution.



a) Do A and B form an ideal solution? Justify your answer.

b) If A and B do not form an ideal solution, are the A--B interactions stronger than, approximately equal to, or weaker than the average of the A---A and B---B interactions? Justify your answer.

c) What is p_B° , the vapor pressure of pure B, at the temperature corresponding to the diagram above?

16) A solution contains a mixture of pentane and hexane at room temperature. The solution has a vapor pressure of 258. torr. Pure pentane and hexane have vapor pressures of 425 torr and 151 torr, respectively, at room temperature. What is the mole fraction composition of the mixture? Assume ideal behavior.

17) How many grams of glucose ($C_6H_{12}O_6$, MW = 180.2 g/mol) must be added to 552. g of water (H_2O , MW = 18.02 g/mol) to give a solution with a vapor pressure 0.20 torr less than that of pure water at 20.0 °C? The vapor pressure of pure water at this temperature is $p_{H_2O}^\circ = 17.5$ torr.

18) An ethylene glycol solution contains 21.2 g of ethylene glycol ($C_2H_6O_2$, MW = 62.07 g/mol) in 85.4 mL of water, (H_2O , MW = 18.02 g/mol). Determine the freezing point and boiling point of the solution. Assume a density of 1.000 g/mL for water. Note that for water $K_f = 1.86$ °C·kg/mol and $K_b = 0.512$ °C·kg/mol, and that ethylene glycol is a nonvolatile and nonionizing solute.

19) An aqueous solution containing 35.9 g of an unknown nonvolatile and nonionizing molecular compound in 150.0 g of water was found to have a freezing point of $-1.30\text{ }^{\circ}\text{C}$. Calculate the molar mass of the unknown compound. For water, $K_f = 1.86\text{ }^{\circ}\text{C}\cdot\text{kg/mol}$.

20) The density of a 0.900 % by mass aqueous solution of sodium chloride (NaCl , $\text{MW} = 58.44\text{ g/mol}$) is $D = 1.0046\text{ g/mL}$ at $T = 25.0\text{ }^{\circ}\text{C}$. What is the osmotic pressure of the above solution (in atm), relative to pure water, at this temperature?

21) What is the osmotic pressure, in atmospheres, of a 1.57 M aqueous solution of urea ($(\text{NH}_2)_2\text{CO}$, $\text{MW} = 60.06\text{ g/mol}$), a nonvolatile and nonionizing solute, at $T = 27.0\text{ }^{\circ}\text{C}$.

22) An isotonic saline solution is an aqueous solution that has approximately the same value for osmotic pressure as human body cells. Isotonic saline contains 9.0 g of sodium chloride (NaCl , $\text{MW} = 58.44\text{ g/mol}$) per liter of solution.

a) What is the molarity of sodium chloride in an isotonic saline solution?

b) What is the osmotic pressure (relative to pure water) of an isotonic saline solution? Assume $T = 37.0\text{ }^{\circ}\text{C}$. Give your final answer in units of atm.

c) Based on your answer to b, explain why red blood cells rupture when placed in deionized water.

23) Arrange the following aqueous solutions in order of decreasing freezing point, and explain your reasoning: 0.50 m HCl , 0.50 m glucose, 0.50 m CH_3COOH .

24) A quantity of 7.480 g of an organic compound is dissolved in water to make 300.0 mL of solution. The solution has an osmotic pressure of 1.43 atm at $T = 27.0\text{ }^{\circ}\text{C}$. The analysis of this compound shows that it contains 41.8 percent C, 4.7 percent H, 37.3 percent O, and 16.3 percent N. Find the molecular formula for the compound.

25) A solution is prepared by dissolving 2.18 g of potassium nitrate (KNO_3 , $\text{MW} = 101.1\text{ g/mol}$) in water. The final volume of the solution is $V = 400.0\text{ mL}$. What is the osmotic pressure of the solution at $T = 20.0\text{ }^{\circ}\text{C}$?

26) Consider 0.0100 mol/kg aqueous solutions of iron III chloride (FeCl_3), calcium nitrate ($\text{Ca}(\text{NO}_3)_2$), and copper II sulfate (CuSO_4). Which of these solutions will have the highest freezing point, and which of these solutions will have the highest melting point? Justify your answer.

27) What is a colloid? How are colloids similar to solutions, and how do they differ from solutions?