

**FORMULA SHEET (tear off)**

1A										8A							
1 H 1.01	2A										3A	4A	5A	6A	7A	2 He 4.00	
3 Li 6.94	4 Be 9.01											5 B 10.81	6 C 12.01	7 N 14.01	8 O 16.00	9 F 19.00	10 Ne 20.18
11 Na 22.99	12 Mg 24.31											13 Al 26.98	14 Si 28.09	15 P 30.97	16 S 32.07	17 Cl 35.45	18 Ar 39.95
19 K 39.10	20 Ca 40.08	21 Sc 44.96	22 Ti 47.87	23 V 50.94	24 Cr 52.00	25 Mn 54.94	26 Fe 55.85	27 Co 58.93	28 Ni 58.69	29 Cu 63.55	30 Zn 65.41	31 Ga 69.72	32 Ge 72.64	33 As 74.92	34 Se 78.96	35 Br 79.90	36 Kr 83.80
37 Rb 85.47	38 Sr 87.62	39 Y 88.91	40 Zr 91.22	41 Nb 92.91	42 Mo 95.94	43 Tc [98]	44 Ru 101.1	45 Rh 102.9	46 Pd 106.4	47 Ag 107.9	48 Cd 112.4	49 In 114.8	50 Sn 118.7	51 Sb 121.8	52 Te 127.6	53 I 126.9	54 Xe 131.3
55 Cs 132.9	56 Ba 137.3	71 Lu 175.0	72 Hf 178.5	73 Ta 181.0	74 W 183.8	75 Re 186.2	76 Os 190.2	77 Ir 192.2	78 Pt 195.1	79 Au 197.0	80 Hg 200.6	81 Tl 204.4	82 Pb 207.2	83 Bi 209.0	84 Po [209]	85 At [210]	86 Rn [222]
87 Fr [223]	88 Ra [226]	103 Lr [262]	104 Rf [261]	105 Db [262]	106 Sg [266]												
		57 La 138.9	58 Ce 140.1	59 Pr 140.9	60 Nd 144.2	61 Pm [145]	62 Sm 150.4	63 Eu 152.0	64 Gd 157.2	65 Tb 158.9	66 Dy 162.5	67 Ho 164.9	68 Er 167.3	69 Tm 168.9	70 Yb 173.0		
		89 Ac [227]	90 Th 232.0	91 Pa 231.0	92 U 238.0	93 Np [237]	94 Pu [244]	95 Am [243]	96 Cm [247]	97 Bk [247]	98 Cf [251]	99 Es [252]	100 Fm [257]	101 Md [258]	102 No [259]		

$$N_A = 6.022 \times 10^{23}$$

$$1 \text{ amu} = 1.661 \times 10^{-27} \text{ kg}$$

$$1 \text{ atm} = 760 \text{ torr} = 760 \text{ mm Hg}$$

$$R = 0.08206 \text{ L}\cdot\text{atm}/\text{mol}\cdot\text{K}$$

$$R = 8.314 \text{ J}/\text{mol}\cdot\text{K}$$

$$^\circ\text{C} = (5/9) (^\circ\text{F} - 32)$$

$$^\circ\text{C} = \text{K} - 273.15$$

$$1 \text{ atm} = 1.013 \text{ bar}$$

$$1 \text{ L}\cdot\text{atm} = 101.3 \text{ J}$$

$$1 \text{ J} = 1 \text{ kg}\cdot\text{m}^2/\text{s}^2$$

$$^\circ\text{F} = (9/5)(^\circ\text{C}) + 32$$

$$\text{K} = ^\circ\text{C} + 273.15$$

$$pV = nRT$$

$$p_A = X_A p_A^\circ$$

$$\Delta T_b = K_b m_B$$

$$H = U + pV$$

$$[B] = k p_B$$

$$\Delta T_f = K_f m_B$$

$$G = H - TS$$

$$\Delta p_A = X_B p_A^\circ$$

$$\Pi = M_B RT$$

**GENERAL CHEMISTRY 2  
FIRST HOUR EXAM  
SEPTEMBER 24, 2021**

**Name** \_\_\_\_\_

**Panthersoft ID** \_\_\_\_\_

**Signature** \_\_\_\_\_

**Part 1** \_\_\_\_\_ (20 points)

**Part 2** \_\_\_\_\_ (42 points)

**Part 3** \_\_\_\_\_ (38 points)

**TOTAL** \_\_\_\_\_ (100 points)

**Do all of the following problems. Show your work.**

**Part 1. Multiple choice.** Circle the letter corresponding to the correct answer. There is one and only one correct answer per problem. [4 points each]

- 1) Which of the following will make it more likely that a mixture of two liquids will form a solution?
  - a) An increase in energy when the liquids form a solution
  - b) An increase in randomness when the liquids form a solution
  - c) A decrease in randomness when the liquids form a solution
  - d) Both a and b
  - e) Both a and c
  
- 2) Which of the following combinations of two liquids is likely to form a solution?
  - a) A combination of two polar liquids
  - b) A combination of two nonpolar liquids
  - c) A combination of a polar liquid and a nonpolar liquid
  - d) Both a and b
  - e) Both b and c
  
- 3) A sample of a metal alloy has a mass of 273.5 g. It contains 473. ppm by mass cobalt. The number of grams of cobalt in the metal alloy is
  - a) 0.0129 g
  - b) 0.129 g
  - c) 1.29 g
  - d) 12.9 g
  - e) 129. g
  
- 4) For a chemical reaction taking place at a constant temperature and standard conditions, which of the following must be true?
  - a)  $\Delta G^\circ_{\text{rxn}} < 0$
  - b)  $\Delta G^\circ_{\text{rxn}} > 0$
  - c)  $\Delta S^\circ_{\text{rxn}} > 0$
  - d) Both a and c
  - e) Both b and c
  
- 5) For a particular chemical reaction the values for  $\Delta H^\circ_{\text{rxn}}$  is positive, and the value for  $\Delta S^\circ_{\text{rxn}}$  is negative. Based on this information we would expect
  - a) The reaction is spontaneous at low temperatures, but not at high temperatures
  - b) The reaction is spontaneous at high temperatures, but not at low temperatures
  - c) The reaction is always spontaneous
  - d) The reaction is never spontaneous
  - e) Any of the above might be expected to be true

**Part 2. Short answer.**

1) Define the following terms [4 points each]

a) heterogeneous mixture –


b) Third Law of Thermodynamics –

2) The figure below is for a mixture of two volatile liquids A and B, at  $T = 40.0\text{ }^{\circ}\text{C}$ . Based on the information in the figure, answer the following questions? [4 points each]

a) Do A and B form an ideal solution (yes or no, and a brief justification for your answer)?

b) What is  $p_B^{\circ}$ , the vapor pressure of pure B? Give your final answer in units of torr.


c) What is  $p_B$ , the partial pressure of B, for a solution with  $X_A = 0.40$ ? Give your final answer in units of torr.



3) A solution is formed by dissolving 13.81 g of glucose ( $C_6H_{12}O_6$ , MW = 180.2 g/mol) in water ( $H_2O$ , MW = 18.02 g/mol). The final volume of the solution is  $V = 250.0$  mL. What is the molarity of glucose in the solution? [8 points]

4) Give the formation reaction for nickel II carbonate ( $NiCO_3(s)$ ). [5 points]

5) Consider the following process (called fusion, or melting) involving an unknown metal M.



The change in enthalpy and the change in entropy for the process, measured for standard conditions and  $T = 25.0$  °C, are  $\Delta H^\circ_{\text{fus}} = 6.68$  kJ/mol, and  $\Delta S^\circ_{\text{fus}} = 9.17$  J/mol·K.

a) Which phase of the metal has a larger value for entropy at standard conditions and  $T = 25.0$  °C, the solid or the liquid (circle the correct answer). [4 points]

M(s) has the larger  
value for  $S^\circ$

M( $\ell$ ) has the larger  
value for  $S^\circ$

b) Estimate the value for  $T^\circ_{\text{fus}}$ , the normal melting point for the metal. Give your final answer in units of °C. [5 points]


**Part 3. Problems.**

1) A solution is prepared by dissolving 30.83 g of a nonvolatile and nonionizing pure chemical compound X in benzene ( $C_6H_6$ , MW = 78.11 g/mol). The final mass of the solution is mass = 605.2 g, and the molality of X in the solution is  $m_X = 0.3627$  mol/kg.

a) Based on this information, find the molecular weight of X. [12 points]

b) The normal freezing point for benzene is  $T_f^\circ = 5.5$  °C, and the freezing point depression constant for benzene is  $K_f = 5.12$  kg·°C/mol. Based on this information and the other information given in the problem, find the freezing point temperature for the solution of X + benzene. Give your final answer in units of °C. [8 points]

2) Thermochemistry is often used to study the behavior of ions in aqueous solution. Consider the following process involving sodium sulfate ( $\text{Na}_2\text{SO}_4$ )



a) What is the value for  $i$ , the van't Hoff factor, for sodium sulfate? Circle the correct answer. [4 points]

$i = 1$

$i = 2$

$i = 3$

$i = 4$

b) Based on the data below, find the value for  $\Delta S^\circ_{\text{rxn}}$  and  $\Delta G^\circ_{\text{rxn}}$  for the above reaction, at  $T = 25.0^\circ\text{C}$ . The data in the table below is also given at  $T = 25.0^\circ\text{C}$ . [10 points]

substance	$\Delta H^\circ_f$ (kJ/mol)	$\Delta G^\circ_f$ (kJ/mol)	$S^\circ$ (J/mol·K)
$\text{Na}^+(\text{aq})$	- 240.1	- 261.9	59.0
$\text{SO}_4^{2-}(\text{aq})$	- 909.3	- 744.5	20.1
$\text{Na}_2\text{SO}_4(\text{s})$	- 1387.1	- 1270.2	149.6

c) Is the above reaction spontaneous for standard conditions and  $T = 25.0^\circ\text{C}$ ? (yes or no, and a brief justification for your answer). [4 points]