

**GENERAL CHEMISTRY 2
SECOND HOUR EXAM (Sample)**

Name _____

Panthersoft ID _____

Signature _____

Part 1 _____ (20 points)

Part 2 _____ (32 points)

Part 3 _____ (48 points)

TOTAL _____ (100 points)

j


Do all of the following problems. Show your work.

Part 1. Multiple choice. Circle the letter corresponding to the correct answer. There is one and only one correct answer per problem. [4 points each]

1) For which of the following reactions can an expression be written for both K_C and K_p ?

- a) $2 \text{H}_2(\text{g}) + \text{C}_2\text{H}_2(\text{g}) \rightleftharpoons \text{C}_2\text{H}_6(\text{g})$
- b) $\text{MgCO}_3(\text{s}) \rightleftharpoons \text{MgO}(\text{s}) + \text{CO}_2(\text{g})$
- c) $\text{HNO}_2(\text{aq}) + \text{H}_2\text{O}(\ell) \rightleftharpoons \text{H}_3\text{O}^+(\text{aq}) + \text{NO}_2^-(\text{aq})$
- d) Both a and b
- e) Both a and b and c

2) Consider the following chemical reaction at $T = 25.^\circ\text{C}$.



The equilibrium constant for this reaction at this temperature is $K_C = 0.0046$

For a closed system with an initial value for reaction quotient $Q_C = 1.00$ at $T = 25.^\circ\text{C}$, which of the following will occur as the system approaches equilibrium?

- a) Moles of N_2O_4 will increase
 - b) Moles of NO_2 will decrease
 - c) Moles of NO_2 will increase
 - d) Both a and b
 - e) Both a and c
- 3) Which of the following compounds is a strong soluble base?
- a) $\text{Ba}(\text{OH})_2$
 - b) $\text{Co}(\text{OH})_2$
 - c) $\text{Fe}(\text{OH})_3$
 - d) Both a and b
 - e) Both a and b and c
- 4) Which of the following substances is classified as a salt?
- a) NaCl
 - b) NaNO_3
 - c) NaOH
 - d) Both a and b
 - e) Both a and b and c

5) In the reaction




the $\text{Al}(\text{H}_2\text{O})_6^{3+}$ ion functions as

- a) the conjugate acid of H_2O
- b) the conjugate base of H_2O
- c) a Bronsted acid
- d) a Bronsted base
- e) none of the above

Part 2. Short answer.

1) A system containing the gases Cl_2 , NO , and NOCl will achieve equilibrium. The process that takes place is



Consider a system initially at equilibrium. For each of the changes given below, indicate (by circling the correct answer) whether the change will lead to an increase, decrease, or no change in the number of moles of Cl_2 in the system. In all cases temperature is held constant at $T = 500. \text{ K}$. [4 points each]

Add 0.20 moles of NOCl to the system under conditions of constant volume

moles of Cl_2
will increase

moles of Cl_2
will remain constant

moles of Cl_2
will decrease

Double the volume of the system

moles of Cl_2
will increase

moles of Cl_2
will remain constant

moles of Cl_2
will decrease


Addition of a catalyst to the system

moles of Cl_2
will increase

moles of Cl_2
will remain constant

moles of Cl_2
will decrease

2) The equilibrium constant for the reaction



is $K_C = 8.1 \times 10^{-24}$ at $T = 25. \text{ }^\circ\text{C}$.

A system initially contains 0.0840 mol/L of $\text{F}_2\text{(g)}$ at $T = 25. \text{ }^\circ\text{C}$. No F(g) is initially present in the system. What will the concentration of F(g) be in the system when equilibrium is reached? [10 points]

3) Define the following [5 points each]

amphoteric


Lewis base

Part 3. Problems.


1) Thermodynamic data are given below (at $T = 25. \text{ }^\circ\text{C}$) and may be of use in doing this problem.

Substance	ΔH°_f (kJ/mol)	ΔG°_f (kJ/mol)	S° (J/mol·K)
$\text{Br}_2(\text{g})$	30.907	3.110	245.46
$\text{Cl}_2(\text{g})$	0.00	0.00	223.07
$\text{CH}_3\text{Br}(\text{g})$	- 35.4	- 26.3	246.4
$\text{CH}_3\text{Cl}(\text{g})$	- 81.9	- 58.4	234.6

What are $\Delta G^\circ_{\text{rxn}}$ and K_{eq} for the following reaction, at $T = 25. \text{ }^\circ\text{C}$ [12 points]



2) The numerical value for the equilibrium constant for the reaction



at $T = 25.^\circ\text{C}$ is $K_C = 7.90$

a) What is the numerical value for K_p for the above reaction at $T = 25.^\circ\text{C}$? [4 points]

b) A system at $T = 25.^\circ\text{C}$ initially has $0.100 \text{ mol/L Cl}_2(\text{g})$ and $0.360 \text{ mol/L BrCl}(\text{g})$. No $\text{Br}_2(\text{g})$ is initially present in the system. What will be the concentrations of BrCl , Br_2 , and Cl_2 when the system reaches equilibrium? [14 points]

3) Find the pH of the following two solutions (both at $T = 25. \text{ }^\circ\text{C}$).

a) 6.44 g of potassium hydroxide (KOH, MW = 56.11 g/mol), a strong soluble base, dissolved in water, to form a solution of final volume $V = 400.0 \text{ mL}$. [8 points]

b) A 0.0800 M solution of propionic acid, a weak monoprotic acid, with $K_a = 1.4 \times 10^{-5}$. [10 points]