

**GENERAL CHEMISTRY 2
FINAL EXAM**

Name _____

Panthersoft ID _____

Signature _____

Part 1 _____ (48 points)

Part 2 _____ (64 points)

Part 3 _____ (88 points)


TOTAL _____ (200 points)

Do all of the following problems. Show your work.

Part 1. Multiple choice. Circle the letter corresponding to the correct answer. There is one and only one correct answer per problem. [4 points each]

- 1) Which of the following methods for expressing concentration have no units?
- molality
 - molarity
 - mole fraction
 - Both a and b
 - All of the above methods for expressing concentration have units
- 2) A solution is formed by adding 0.100 moles of a nonvolatile and nonionizing solute to 250.0 g of a volatile solvent. Compared to the pure solvent
- the boiling point of the solution will be higher than the boiling point of the pure solvent
 - the freezing point of the solution will be higher than the freezing point of the pure solvent
 - the vapor pressure of the solution will be higher than the vapor pressure of the pure solvent
 - Both a and b
 - Both b and c
- 3) For a spontaneous reaction which of the following must be true?
- $\Delta G_{\text{rxn}} < 0$
 - $\Delta S_{\text{rxn}} < 0$
 - $\Delta S_{\text{rxn}} > 0$
 - Both a and b
 - Both a and c
- 4) For which of the following reactions would we expect $\Delta S^{\circ}_{\text{rxn}}$ to be large and negative?
- $\text{MgCO}_3(\text{s}) \rightarrow \text{MgO}(\text{s}) + \text{CO}_2(\text{g})$
 - $\text{C}_6\text{H}_6(\ell) \rightarrow \text{C}_6\text{H}_6(\text{g})$
 - $\text{C}_3\text{H}_8(\text{g}) + 5 \text{O}_2(\text{g}) \rightarrow 3 \text{CO}_2(\text{g}) + 4 \text{H}_2\text{O}(\ell)$
 - both a and b
 - both a and c

5) Consider the reaction




A system containing I_2 , Cl_2 , and ICl is initially at equilibrium at some temperature T . The volume of the system is decreased to half of its initial value while keeping temperature constant. Which of the following will occur as the system re-establishes equilibrium?

- The moles of I_2 in the system will increase
 - The moles of Cl_2 in the system will increase
 - The moles of ICl in the system will increase
 - Both a and b
 - None of the above
- 6) Benzoic acid ($\text{C}_6\text{H}_5\text{COOH}$) is a weak acid, with $K_a = 6.5 \times 10^{-5}$ (at $T = 25.^\circ\text{C}$). The $\text{C}_6\text{H}_5\text{COO}^-$ ion is
- a strong acid
 - a weak acid
 - a strong base
 - a weak base
 - None of the above, as ions have no acid or base properties

- 7) A Bronsted acid is
- a proton acceptor
 - a proton donor
 - an electron pair acceptor
 - an electron pair donor
 - both b and c
- 8) Consider the following three substances: HBr, H₂S, and H₂Se. Of these substances
- HBr is the strongest acid and H₂S is the weakest acid
 - HBr is the strongest acid and H₂Se is the weakest acid
 - H₂S is the strongest acid and HBr is the weakest acid
 - H₂Se is the strongest acid and HBr is the weakest acid
 - H₂Se is the strongest acid and H₂S is the weakest acid
- 9) Which of the following aqueous solutions will be a buffer solution?
- A solution containing 0.100 M HBr (an acid) and 0.050 M NaBr.
 - A solution containing 0.100 M HNO₂ (an acid) and 0.050 M NaNO₂.
 - A solution containing 0.100 M HNO₂ (an acid) and 0.050 M NaOH.
 - Both a and b
 - Both b and c

- 10) The standard reduction potential for the process



is $E^{\circ} = + 0.34 \text{ v}$. Based on this, we can say that the half-cell oxidation potential for the process



is

- $E^{\circ} = + 0.68 \text{ v}$
 - $E^{\circ} = + 0.34 \text{ v}$
 - $E^{\circ} = 0.00 \text{ v}$
 - $E^{\circ} = - 0.34 \text{ v}$
 - $E^{\circ} = - 0.68 \text{ v}$
- 11) A chemical reaction obeys the rate law

$$\text{rate} = k [\text{A}] [\text{B}]^2$$

The overall order of the reaction is

- first order
 - second order
 - third order
 - both a and b
 - both a and b and c
- 12) A catalyst
- changes the rate of reaction and also changes the equilibrium constant for a reaction
 - changes the rate of reaction but does not change the equilibrium constant for a reaction
 - does not change the rate of reaction but changes the equilibrium constant for a reaction
 - does not change the rate of reaction and also does not change the equilibrium constant for a reaction
 - cannot be present in a system at equilibrium

Part 2. Short answer.

1) Define the following terms [6 points each]

a) amphoteric –

b) catalyst –

2) 0.45 g of a nonvolatile pure chemical substance (a polymer) are dissolved in liquid benzene (C_6H_6 , MW = 78.1 g/mol), to form a solution with final volume $V = 200.0$ mL. The osmotic pressure of the solution, measured at $T = 20.0$ °C, was 33.2 torr. Based on this information find the molecular weight of the polymer. [10 points]

3) For each of the following questions circle the correct answer. There is one and only one correct answer per problem. [4 points each]

a) The substance with $\Delta G^\circ_f = 0.0$ kJ/mol at $T = 25$. °C

$CuCl_2(s)$

$C_6H_{12}(\ell)$

$O_2(g)$

$O_3(g)$

b) The acid that is a polyprotic acid

HBr

$HBrO_2$

HNO_3

H_3PO_4

c) The hydroxide compound that is a strong soluble base

AgOH

$Ba(OH)_2$

$Cu(OH)_2$

$Fe(OH)_3$

d) The best indicator to use in the titration of a strong acid by a strong base

alizarin yellow
 $pK_{ind} = 11.0$

bromothymol blue
 $pK_{ind} = 6.8$

bromophenol blue
 $pK_{ind} = 3.8$

thymol blue
 $pK_{ind} = 2.0$


4) Equilibrium between sulfur oxides in the presence of molecular oxygen is difficult to study experimentally. Consider the following reaction



a) Give the expression for K_C , the equilibrium constant, in terms of reactant and product concentrations. [4 points]

b) For a particular system the initial concentrations of SO_3 and O_2 are both 0.0400 mol/L. No SO_2 is initially present. Find the value for $[\text{SO}_2]$, the concentration of SO_2 , when equilibrium is reached. [10 points]

5) Balance the following oxidation-reduction reaction for the indicated condition. [12 points each]




Part 3. Problems.


1) Thermochemical data (at $T = 25. \text{ }^\circ\text{C}$) for several substances is given below, and may be of use in doing the following problems, which also take place at $T = 25. \text{ }^\circ\text{C}$.

Substance	ΔH°_f (kJ/mol)	ΔG°_f (kJ/mol)	S° (J/mol·K)
C(s)	0.00	0.00	5.74
CO ₂ (g)	- 393.51	- 394.36	213.74
CO(NH ₂) ₂ (s)	- 333.51	- 197.33	104.60
H ₂ O(g)	- 241.82	- 228.57	188.83
NH ₃ (g)	- 46.11	- 16.45	192.45
O ₂ (g)	0.00	0.00	205.14

a) What is the value for $\Delta S^\circ_{\text{rxn}}$ for the reaction [8 points]



b) What are the values for $\Delta G^\circ_{\text{rxn}}$ and K for the reaction [16 points]



2) Titration is a common method for finding the concentration of stock solutions of acids or bases. The following problem concerns a titration carried out at $T = 25. \text{ }^{\circ}\text{C}$.

A student titrates a 25.00 mL sample of a 0.1814 M solution of hypochlorous acid (HClO , $K_a = 2.9 \times 10^{-8}$) with a stock solution of sodium hydroxide (KOH), a strong soluble base.

a) What is the initial pH of the sample HClO solution? [8 points]





b) After 38.17 mL of sodium hydroxide solution is added the equivalence point for the titration is reached. Based on this result, what is the concentration of the sodium hydroxide stock solution? [8 points]

c) In the above titration, what is the pH at the equivalence point (when 25.00 mL of stock HClO solution has been mixed with 38.17 mL of stock NaOH solution)? [12 points]

3) Consider the following galvanic cell (at $T = 25. \text{ }^\circ\text{C}$)



Find the following. You may need to use some of the reduction data given below in doing part of this problem



a) The half cell oxidation reaction, the half cell reduction, and the net cell reaction [5 points]

b) E°_{cell} [5 points]

c) E_{cell} [6 points]

