

CHM 3411 – Problem Set 2

Due date: Friday, January 25<sup>th</sup> (I've pushed the due date back due to the MLK holiday on Monday)

Do all of the following problems. Show your work.

1) For each of the following operators and functions, determine under what conditions (if any) the function is an eigenfunction of the operator. For those functions that are eigenfunctions, find the corresponding eigenvalues.

a)  $\hat{O} = d^2/dx^2$  ,  $f(x) = \exp(ax)$  , where  $a$  is a positive constant

b)  $\hat{O} = d^2/dx^2$  ,  $f(x) = \cos(ax)$  , where  $a$  is a positive constant

c)  $\hat{O} = x (d/dy) + y (d/dx)$  ,  $f(x,y) = x^2 + y^2$

d)  $\hat{O} = x (d/dy) + y (d/dx)$  ,  $f(x,y) = xy$

e)  $\hat{O} = \text{inv}$  ,  $f(x) = x^2 + x$  (The inversion operator,  $\text{inv}$ , replaces  $x$  with  $-x$  everywhere it appears)

f)  $\hat{O} = \text{inv}$  ,  $f(x) = x^2 + 1$

2) Consider the following function

$$f(x) = Nx^2 \quad -L \leq x \leq L, L > 0$$

$$f(x) = 0 \quad x < -L \text{ or } x > L$$

a) Find the value for  $N$  that makes  $f(x)$  a normalized function.

b) Would  $f(x)$  be an acceptable solution to some TISE? Why or why not? (Hint: Does  $f(x)$  satisfy all of the requirements for a wavefunction?)

3) There are a number of proofs in quantum mechanics that are both useful and surprisingly easy. Here we explore one of these.

Let  $f(x)$  be a “well behaved” normalized function. Because the functions  $\psi_n$  that are solutions to a particular TISE form a complete orthonormal set of functions (basis functions), we can say:

$$\begin{aligned} f(x) &= c_1 \psi_1(x) + c_2 \psi_2(x) + c_3 \psi_3(x) + \dots \\ &= \sum_{n=1}^{\infty} c_n \psi_n(x) \end{aligned} \quad (3.1)$$

a) Left multiply the left and right sides of eq 3.1 by  $\psi_j^*$ , the complex conjugate of the  $j^{\text{th}}$  solution to the TISE.

b) Integrate all of the terms from a over the region  $-\infty < x < \infty$ .

c) Using the orthonormality condition, find an expression for  $c_j$ , the coefficient for the  $j^{\text{th}}$  term in eq 3.1.

d) The result in part c takes on a particularly meaningful form if  $f(x)$  is a normalized function, that is, if

$$\int_{-\infty}^{\infty} f(x)^* f(x) dx = 1 \quad (3.2)$$

Using your result from part c, show that if  $f(x)$  is a normalized function, it follows that

$$|c_1|^2 + |c_2|^2 + |c_3|^2 + \dots = \sum_{n=1}^{\infty} |c_n|^2 = 1 \quad (3.3)$$

where

$$|c_n|^2 = c_n^* c_n \quad (3.4)$$

(Hint: Use the normalization condition (eq 3.2) as a starting point.)

The significance of this result is that if  $f(x)$  is the wavefunction for a state in a system, and if  $f(x)$  is a normalized function, then  $|c_n|^2$  represents the probability of being in the  $\psi_n^{\text{th}}$  eigenstate of the system.